

# Oviposition habitat selection for a predator refuge and food source in a mosquito

J. GUILLERMO BOND<sup>1,2</sup>, JUAN I. ARREDONDO-JIMÉNEZ<sup>1</sup>, MARIO H. RODRÍGUEZ<sup>3</sup>, HUMBERTO QUIROZ-MARTÍNEZ<sup>4</sup> and TREVOR WILLIAMS<sup>2,5</sup>  
<sup>1</sup>Centro de Investigación de Paludismo-INSP, Tapachula, Mexico, <sup>2</sup>ECOSUR, Tapachula, Mexico, <sup>3</sup>Centro de Investigaciones sobre Enfermedades Infecciosas-INSP, Cuernavaca, Mexico, <sup>4</sup>Facultad de Ciencias Biológicas, Universidad Autónoma Nuevo León, San Nicolás de los Garza, Mexico and <sup>5</sup>Departamento de Producción Agraria, Universidad Pública de Navarra, Pamplona, Spain

**Abstract.** 1. The influence of filamentous algae on oviposition habitat selection by the mosquito *Anopheles pseudopunctipennis* and the consequences of oviposition decisions on the diet, development, body size, and survival of offspring were examined.

2. A natural population of *An. pseudopunctipennis* in Chiapas, Mexico, oviposited almost exclusively in containers with filamentous algae. Algae represented 47% of the gut contents of mosquito larvae sampled from the natural population. Mosquito larvae fed on an exclusive diet of algae developed as quickly and achieved the same adult body size (wing length) as their conspecifics fed on a standard laboratory diet.

3. Multiple regression of survival of mosquito larvae on percentage surface area cover of algae (0–99%) and the density of predatory fish (zero to four fish per container) was best described by a second-order polynomial model. Increasing fish densities resulted in a reduction in mosquito survival in all algal treatments. The highest incidence of survival was observed at intermediate (66%) algal cover in all treatments.

4. The presence of fish significantly extended larval development times whereas algal cover had no significant effect. The presence of fish resulted in emergence of smaller adults due to reduced feeding opportunities and predator avoidance behaviour. Algal cover also affected mosquito wing length but differently at each fish density.

5. Oviposition habitat selection improves survival in the presence of predators and feeding opportunities for *An. pseudopunctipennis* larvae.

**Key words.** *Anopheles pseudopunctipennis*, *Cladophora glomerata*, feeding, fish, gut analysis, offspring survival, oviposition, predators, *Spirogyra majuscula*.

## Introduction

Ovipositing insects should select sites that improve the survival, growth, and reproductive potential of the

offspring, especially for species in which juveniles are incapable of migrating away from low quality habitats (Peckarsky *et al.*, 2000). In the absence of inter- or intra-specific competition, progeny fitness will depend on two factors: an adequate food supply and the presence of natural enemies (predators, parasites, and pathogens). As a result, insects often avoid ovipositing in sites where there is a high risk of predation to their progeny (Blaustein, 1999; Angelon & Petranka, 2002; Spencer *et al.*, 2002; Kiflawi

Correspondence: Trevor Williams, Departamento de Producción Agraria, Universidad Pública de Navarra, Pamplona 31006, Spain. E-mail: trevor.williams@unavarra.es

et al., 2003). Oviposition habitat selection is particularly relevant in insect vectors of medical importance as it determines the localities to which larvicidal control measures are targeted (Service, 1993).

Understanding the factors that regulate the size of mosquito populations is considered fundamental to the ability to predict transmission rates and for vector population control (Service, 1989, 1995). Aquatic vegetation can exert an important influence on the reproduction of mosquitoes (Orr & Resh, 1989, 1992; Kant et al., 1996). This may involve facilitating oviposition (Orr & Resh, 1992; Rejmankova et al., 1996) or by providing a larval microhabitat (Orr & Resh, 1992) from which to escape the attention of predators (Sih, 1986; Orr & Resh, 1989). Mosquitoes may also select oviposition sites based on the availability of larval food (Blaustein & Kotler, 1993). Studies of the factors that determine anopheline larval population densities have almost invariably assumed that aquatic vegetation represents an important food resource for larvae (Hall, 1972; Rejmankova et al., 1992; Marten et al., 1996), although anopheline feeding on filamentous algae is poorly studied.

Environmental management based on ecological principals represents an effective and expedient intervention for insect vector control (Ault, 1994). For example, a direct linear relationship between anopheline larval densities and percentage of breeding pool cover by filamentous algae in southern Mexico has recently been reported (Bond et al., 2004b). Manual extraction of algae from anopheline breeding pools resulted in a marked reduction in larval and adult population densities that was sustained for approximately 2 months, by which time seasonal rains began to physically remove larvae from breeding sites.

The mosquito *Anopheles pseudopunctipennis* Theobald (Diptera: Culicidae) is an important vector of *Plasmodium* spp. in tropical regions of the Americas (Fernández-Salas et al., 1994a,b). An intimate association between filamentous algae and immature stages of *An. pseudopunctipennis* has been reported throughout the geographic range of this mosquito (Rejmankova et al., 1991, 1993; Manguin et al., 1996). The present study was designed to examine the influence of filamentous algae on oviposition habitat selection by *An. pseudopunctipennis* and the consequences of oviposition decisions on the diet, development, body size, and survival of offspring in the presence of predators.

## Methods

### Study site

Field studies were performed in a large concrete water storage pool 20 m × 20 m × 1.5 m deep with water, in the village of El Plan – Las Palmas, Municipality of Tapachula Chiapas, Mexico (15°03'81"N, 92°12'40"W), at an altitude of 315 m above sea level. The pool was continuously fed by a small stream, a tributary of the River Coatán, which flows past the village at a distance of ≈ 300 m

from the water storage pool. Temporary riverside pools along the River Coatán are primary breeding sites of *An. pseudopunctipennis* during the dry season. The climate in this region is tropical sub-humid, with two well-defined seasons: a rainy season from May to October and a dry season from November through April. The annual average rainfall is 3800 mm, the average annual temperature is 25 °C, and the relative humidity is 70–90% for most of the year. Field studies were performed from November 1996 to July 1997.

### Influence of filamentous algae on oviposition habitat selection

Experimental units consisted of PVC tubes (25 cm diameter and 37.5 cm high) with two windows on the sides (5 cm high × 20 cm wide) and a polystyrene foam float glued to the sides of the tube. A fine nylon mesh was glued to the bottom and over the side windows to allow the flow of water and to prevent the escape of mosquito larvae. Ten tubes were tied to a string and fastened to the sides of the pool and submerged so that 10 cm of the tube remained above the water level. Five of the tubes were treated with algae and the remaining five tubes were left as untreated controls. Treatment with algae consisted of placing 170 g (wet weight) of *Spirogyra majuscula* Kützing in each tube to give approximately 66% cover of the water surface. Tubes were collected 3 days later and the number of *An. pseudopunctipennis* larvae was recorded by microscopic examination of the contents of each tube in a field laboratory. The algae from each tube were placed in trays covered with muslin and incubated in the field laboratory for a further 7 days to permit hatching of all viable eggs. An identical procedure was performed using *Cladophora glomerata* (L.). Overall, each species of algae was tested five times in an alternating sequence. The absence of oviposition in tubes lacking algae prevented statistical analysis; the total number of first-instar larvae observed in each replicate (data pooled for all five tubes) was compared for each species of alga by Mann–Whitney *U*-test using the SPSS v.10.0 package (SPSS, 1995).

### Role of filamentous algae in the diet and development of larvae

(i) *Analysis of larval gut contents.* Fourth-instar *An. pseudopunctipennis* larvae were collected from 80 breeding pools (20 larvae per pool), randomly selected within a 6-km section of the River Coatán. Larvae were transported to the laboratory in a cool-box and killed by brief immersion in hot water. The food bolus enclosed in the peritrophic membrane was removed from each larval gut by dissection and placed in groups of 20 in plastic vials with 2 ml of sterile water. Vials were briefly sonicated in an ultrasonic bath to disrupt aggregates. Samples were then stained with DAPI (4',6-diamidino-2-phenylindole) (Sigma, St Louis, Missouri) at a final concentration of 2 µg ml<sup>-1</sup>, and placed in darkness at 4 °C for 25 min. The contents of each vial

were then passed through a 25-mm diameter black polycarbonate filter (Millipore, Bedford, Massachusetts) of 0.45 µm pore size. Each black filter was placed on an oiled microscope slide, covered with immersion oil and a coverslip and examined using a Reichert–Jung Diastar epifluorescence microscope with a ×40 objective. Stained particles were counted in 50 randomly chosen fields of view for each slide. Particles could be differentiated into four categories by their staining: (i) bacteria (blue), (ii) protozoa (blue with distinctive nuclei), (iii) organic detritus (yellow), and (iv) algae (chlorophyll stains red). The number of debris particles, algal fragments and bacterial cells per gut were quantified by counting 50 fields of view, each of 40 × 40 µm area, from each polycarbonate filter following the method of Walker *et al.* (1988) but with no dilution factor correction. The process was repeated for each of the groups of larvae from each of the pools sampled.

(ii) *Suitability of Spirogyra majuscula for larval development.* Groups of 50 first-instar *An. pseudopunctipennis* larvae were obtained from a continuously reared laboratory colony maintained in the Centro de Investigación de Paludismo, Tapachula, Chiapas, Mexico (Villarreal *et al.*, 1998). These larvae hatched within the previous 24 h and were transferred to plastic trays (25 × 15 × 5 cm) with 250 ml of distilled water. Larvae were offered either 1250 mg day<sup>-1</sup> *S. majuscula* or a standard laboratory diet offered at the quantities used for rearing the laboratory colony (30 mg day<sup>-1</sup> during the first and second instars and 90–120 mg day<sup>-1</sup> during the third and fourth instars). To eliminate bacteria, algae were subjected to an antibiotic treatment involving 50 U ml<sup>-1</sup> penicillin G and 50 µg ml<sup>-1</sup> streptomycin sulphate for 48 h followed by extensive washes prior to use in experiments. The standard laboratory colony diet consisted of ground rodent food, mainly comprising ground cereals and soya (23% protein, 6.0% oil, 60% fibre, 4.9% sucrose, 6.1% mineral salts, by weight). Larvae were reared in the laboratory at 28 ± 2 °C and L:D 14:10 h photoperiod. Trays were checked daily and pupae were transferred to water filled plastic cups in emergence cages. The immature development time, from being placed in rearing trays to adult eclosion was recorded. The number and sex of emerging adults was noted daily. Following emergence, the wing length of each mosquito was measured from the axial incision to the apical margin. The fimbrial scales were not included in this measurement and the wing was selected from the left or right side of the body at random. Wing length was taken as an indicator of body size; the cubic function (a dimensionless number) was calculated for statistical analysis as recommended by Briegel (1990a,b). The experiment was performed five times. Results were subjected to *t*-tests with equal and unequal variances, as required.

#### *Filamentous algae as a predator refuge*

(i) *Survival in the presence of predators.* The effects of black molly, *Poecilia sphenops* Valenciennes, a fish

predator of mosquito larvae, and filamentous algae cover on the survival of *An. pseudopunctipennis* larvae were examined in a crossed-design experiment involving four levels of filamentous algae cover (0, 33, 66 and 99%) and four densities of *P. sphenops* (zero, one, two, and four fish per container). Each treatment was replicated four times.

Samples of *Spirogyra* spp. were transported from the River Coatán and were carefully washed and examined to eliminate the presence of predators and other invertebrates. Algae were placed in perforated metallic trays sealed with nylon mesh within the pool, and allowed to establish and recover from the manipulation process for 1 week. Experimental containers consisted of PVC tubes (described above) and were floated with 20 cm of the tube above the water line to prevent fish jumping into or out of the tubes. Sixteen tubes were placed in the water storage pool in groups of four and were held in place by a string fastened to the edge of the pool. Once established, 0, 80, 170, and 250 g algae (wet weight) were added to tubes resulting in 0, 33, 66, or 99% surface cover within the tubes respectively. These quantities of algae were assigned randomly to the experimental tubes and were left to establish for 3 days.

Medium-sized examples of *P. sphenops*, 4–6 cm in total length, were assigned randomly to PVC tubes at densities of zero, one, two, and four fish per tube. Fish were allowed to acclimatise for 24 h. Once acclimatised, 100 first-instar *An. pseudopunctipennis* from the laboratory colony, 24–48 h post-hatching, were added to each container. The number of insects that survived to pupation in each container was recorded daily. The absence of mosquito survival in the treatments involving fish with no algae present resulted in an unbalanced design and precluded a standard factorial analysis of variance. Consequently, the percentage of insects that survived to pupation was  $\sqrt{(x + 1)}$  transformed to normalise the distribution and subjected to quadratic linear regression against the percentage of algal surface cover and fish density. Both algal cover and fish densities were subjected to  $(x - \text{mean } x, y - \text{mean } y)$ , respectively) transformation to reduce collinearity of standard and squared terms in the regression (Sokal & Rohlf, 1981).

(ii) *Effects of algae and predators on development time and adult wing length.* Pupae obtained from each treatment were transported daily to the laboratory in plastic containers in an insulated box. Pupae were placed in plastic cups in mesh cages to permit adult emergence. The number and sex of emerging mosquitoes was noted daily. Adult wing length of 10–30 adults of each sex (depending on survival in each treatment) were measured from each replicate, from the axial incision to the outer margin excluding the fimbrial scales (Armbruster & Hutchinson, 2002). Mean larval development time (the interval from first instar to pupation) and mean adult wing length data were calculated for each replicate and subjected to two-way ANOVA. Mean separation was performed by Tukey test (SPSS, 1995).

## Results

### *Influence of filamentous algae on oviposition habitat selection*

When tested with *S. majuscula*, mosquitoes oviposited exclusively in containers with algae; the mean ( $\pm$  SE) number of first-instar *An. pseudopunctipennis* was  $21.0 \pm 11.4$  larvae per container. No larvae were observed in containers containing water alone. Similarly, when offered *C. glomerata*, oviposition was observed in the treatment involving algae in four out of five replicates (mean of all replicates was  $16.2 \pm 7.2$  larvae per container), whereas larvae were observed in only one replicate (a single oviposition container) in which no algae were present. As a result, the influence of filamentous algae on larval feeding and development and survival in the presence of predators was then studied. A non-parametric comparison between the number of larvae observed in containers with *S. majuscula* and *C. glomerata* indicated no significant difference in the attractiveness of these two species of algae for oviposition of *An. pseudopunctipennis* (Mann–Whitney  $U = 18$ ,  $P > 0.05$ ).

### *Role of filamentous algae in the diet and development of larvae*

(i) *Analysis of larval gut contents.* Of the 80 groups of larvae (20 larvae per group) subjected to gut content analysis, all were positive for the presence of filamentous algae. The mean ( $\pm$  SE) number of particles observed for each group of 20 larval guts was  $70\,180 \pm 2348$  fragments of organic debris,  $64\,197 \pm 2446$  fragments of algae, and  $1227 \pm 276$  bacterial cells. No protozoa were observed. In total, algae represented 47% of the gut contents of mosquito larvae, suggesting that algae comprise an important part of the diet in all the riverside pools from which larvae were sampled.

(ii) *Suitability of Spirogyra majuscula for larval development.* Larval diet had no significant effect on mean wing length (length<sup>3</sup>  $\pm$  SD) of adult females (control =  $58.74 \pm 0.47$  mm; algae =  $58.24 \pm 0.09$  mm;  $t_8 = 0.941$ ,  $P = 0.077$ , unequal variances), or adult males (control =  $45.06 \pm 0.29$  mm; algae =  $46.70 \pm 2.54$  mm;  $t_8 = 0.830$ ,  $P = 0.222$ , unequal variances) in the laboratory. Immature development times were also similar for insects reared on each type of diet for both females (control =  $8.23 \pm 0.10$  days; algae =  $8.19 \pm 0.08$  days;  $t_8 = 0.623$ ,  $P = 0.511$ , equal variances) and males (control =  $8.55 \pm 0.11$  days; algae =  $8.52 \pm 0.24$  days;  $t_8 = 0.443$ ,  $P = 0.807$ , equal variances), indicating that *S. majuscula* and the standard diet used to continuously maintain the laboratory colony were equally suitable for mosquito development.

### *Filamentous algae as a predator refuge*

(i) *Survival in the presence of predators.* Multiple regression of survival of mosquito larvae [expressed as the square

root of the percentage ( $x + 1$ ) of larvae that survived to pupate] on percentage algal cover and fish density revealed a highly significant relationship ( $F_{4,59} = 49.66$ ,  $P < 0.0001$ ; Table 1). The minimal significant model was a second-order polynomial with no interactions between variables.

Increasing fish densities resulted in a reduction in mosquito survival in all algal treatments (Fig. 1). The highest incidence of survival was observed at 66% algal cover in all treatments, such that the relationship between mosquito survival and algal cover was curvilinear at all fish densities (zero to four fish). In the control treatment without fish, mean survival increased from 29% in the absence of algae to 53% at 66% algal cover and then fell to 35% at maximal algal cover (99%). In the presence of fish, mean survival increased from 0 in the absence of algae to maximum values of between 9.0 and 19.5%, in the presence of 66% algal cover, at densities of one and four fish per tube respectively, and then fell slightly (to 5.5 and 16% survival respectively) at the highest (99%) algal cover. The density of two fish per tube was intermediate in all treatments. None of the fish died or escaped during the experimental period.

(ii) *Effects of algae and predators on development time and adult wing length.* The absence of surviving insects in fish treatments without algal cover precluded the determination of development time. These treatments were therefore excluded from the analysis. Development time of larvae, from being released to pupation, was not significantly affected by the percentage of algal cover in any treatment ( $F_{3,39} = 0.017$ ;  $P = 0.974$ ) (Fig. 2). The presence of fish, however, resulted in a significantly extended larval development time ( $F_{3,39} = 42.5$ ;  $P < 0.001$ ), that increased steadily from an average of 8.4 days in treatments without fish, to a maximum of 12.1 days in treatments involving a density of four fish (mean values pooled for 33, 66, and 99% algal treatments within each fish density).

The cube of adult female wing length in millimetres (mean  $\pm$  SD) was  $58.15 \pm 0.16$  in mosquitoes that emerged from the treatment involving no algal cover and no fish. The absence of surviving insects in fish treatments without algal cover precluded measurement of wing length. These treatments were therefore excluded from the analysis. The presence of fish resulted in emergence of adult females with consistently shorter wings compared with those that developed in the no-fish treatments ( $F_{3,46} = 11048.5$ ;  $P < 0.0001$ ) (Fig. 3). Algal cover also affected female mosquito wing length ( $F_{2,46} = 773.6$ ;  $P < 0.0001$ ) but differently at each fish density reflected in a significant interaction effect ( $F_{6,46} = 538.4$ ;  $P < 0.0001$ ). Algal cover at 66 and 99% resulted in a steady decrease in wing length with increasing fish density. At 33% cover, however, wing length was lowest at the densities of one and two fish but increased slightly but significantly at a density of four fish. A similar pattern was observed between treatments in the wing lengths of adult males (results not shown).

**Table 1.** Multiple linear regression (quadratic model) of square-root transformed percentage survival of mosquito larvae on percentage of surface cover of filamentous algae ( $x - \text{mean}$ ) and number of predatory fish per container ( $y - \text{mean}$ ).

Function =  $3.64906 - 1.21921x + 0.02283y + 0.44572x^2 - 0.000843355y^2$   
 No. of cases = 64, adjusted  $r^2 = 0.7622$

Source	Sums of squares	d.f.	Mean	F	P
Regression	261.12	4	65.28	49.66	<0.0001
Residual	77.56	59	1.31		
Total	338.69				

Term	Coefficient	SE	t(59)	P
Intercept	3.64906	0.289675	12.5971	<0.0001
x (algae)	-1.21921	0.109960	-11.0878	<0.0001
y (fish)	0.02283	0.003885	5.8763	<0.0001
$x^2$	0.44572	0.080844	5.5134	<0.0001
$y^2$	-0.00084	0.000132	-6.4081	<0.0001


Both  $x$  and  $y$  were transformed to  $(x - \text{mean})$  and  $(y - \text{mean})$  to reduce colinearity between normal and squared terms in regression.

## Discussion


Oviposition habitat selection is influenced by a diversity of chemical, physical, and physiological factors. Once attracted to the oviposition site, gravid females use visual (colour, texture, brightness), and olfactory cues (semiochemicals) to decide the suitability of a potential habitat for egg laying (Bentley & Day, 1989). However, it is

ovipositional decision making, modulated by the detection of semiochemicals indicating the presence of predators (Stav *et al.*, 1999; Kiflawi *et al.*, 2003; Eitam & Blaustein, 2004; Blaustein *et al.*, 2004), or immature mosquitoes (Bentley *et al.*, 1976; Laurence & Pickett, 1985; Zahiri & Rau, 1998; Reiskind & Wilson, 2004), including oviposition pheromones (Laurence & Pickett, 1985; Mboera *et al.*, 1999), that has attracted most attention from insect ecologists. In this study, an absolute preference by a natural population of *An. pseudopunctipennis* to oviposit in containers with *S. majuscula* over containers with water alone was observed. This behaviour was almost identical when offered a choice of containers with open water or *C. glomerata*. The consequences of the ovipositional decision to the diet, development rate, and body size of offspring, and the role of filamentous algae as a refuge from predatory fish was then examined.

Anophelines usually feed on microorganisms and detritus (Merritt *et al.*, 1992). Algae may also be common components of the gut contents (Howland, 1930), although their nutritional value, and specifically their digestibility, has been questioned (Laird, 1988). The degree of selectivity of mosquito larval feeding is also a subject of debate with various authors suggesting that gut contents largely reflect the size of the larvae, method of feeding and the availability of different food items in the larval habitat (Laird, 1988; Clements, 1992). A number of studies have indicated that bacterial cells are a major constituent of the larval diet (Walker *et al.*, 1988; Merritt *et al.*, 1990; Rejmankova *et al.*, 2000), although bacteria only represented a minor component of the diet of *An. pseudopunctipennis* in this study. Particle analysis of gut contents does not take into account colloidal or dissolved foodstuffs, although such materials may be important to larval nutrition (Wotton, 1990). Indeed, algal cells may produce nutritionally important exudates that are not apparent by gut particle analysis (Wotton, 1990). In addition, particle counts did not allow



**Fig. 1.** Graphical representation of multiple linear regression of the square root of percentage survival of *Anopheles pseudopunctipennis* larvae, of percentage surface cover of *Spirogyra majuscula* ( $x - \text{mean}$ ), and density of predatory fish ( $y - \text{mean}$ ).




**Fig. 2.** Mean development time (first instar to adult eclosion) of *Anopheles pseudopunctipennis* larvae exposed to 0–99% surface cover of filamentous algae and densities of zero to four predatory fish per tube. Columns headed by identical letters are not significantly different (ANOVA, Tukey,  $P > 0.05$ ). As no mosquitoes survived in the 0% algal cover treatment in the presence of fish, this was excluded from the analysis.

us to estimate the biovolume of ingested matter, which may be a more accurate measure of the importance of each constituent of gut contents (Wetzel & Likens, 1991).

These findings demonstrated that filamentous algae were abundant in the guts of larval *An. pseudopunctipennis* in natural breeding pools. Moreover, laboratory-reared larvae developed as quickly and attained the same adult size (as judged by wing length) given a diet of *S. majuscula* or the standard diet of the laboratory colony, indicating that filamentous algae alone are nutritionally suitable to support the entire development of *An. pseudopunctipennis* larvae. Several other aquatic insects have been shown to be capable of development on a diet of filamentous algae monocultures including ephydrid flies (Zack & Foote, 1978) and Trichoptera (Keiper, 2002).

In the absence of predators, the highest survival of larvae was observed at 66% algal cover. Greater quantities of algae reduced survival possibly by presenting a physical barrier to larval respiration or feeding activity close to the air – water interface. Lesser quantities of algae may have reduced mosquito survival by exposing larvae to harmful levels of ultraviolet light (Donahue & Schindler, 1998; Buma *et al.*, 2003).

The black molly (*P. sphenops*) proved to be a voracious predator of *Anopheles* larvae, completely eliminating them from treatments in which no algal refuge was available. However, filamentous algae functioned as an effective refuge from fish predation at densities of one to four fish per container. Survival was highest in containers with 66% algal cover in all treatments involving fish. Submerged



**Fig. 3.** Effects of algal cover (*Spirogyra* spp.) and fish density (*Poecilia sphenops*) on cube of adult female wing length in millimetres (mean ± SD) of *Anopheles pseudopunctipennis*. Columns headed with different capital letters are significantly different (ANOVA, Tukey,  $P < 0.001$ ) for comparisons between fish densities within each algal treatment, columns headed with different lower case letters are significantly different (ANOVA, Tukey,  $P < 0.001$ ) for comparisons between algal cover treatments within each fish density. Larval survival was zero in the treatments involving no algal cover in the presence of fish (excluded from analysis).

vegetation impedes the movement and obstructs the vision of predatory fish (Manatunge *et al.*, 2000) and the density and architectural complexity of the macrophytes are important in that respect (Savino & Stein, 1989; Persson & Eklov, 1995). It was observed that mats of filamentous algae were almost impenetrable to *P. sphenops* limiting them to foraging at the edges of the algal mass.

The results of this study resemble those of Orr and Resh (1989, 1992) who reported a positive relationship between the density of emergent stems in monospecific beds of *Myriophyllum aquaticum* (Vell.) and the density of eggs and larvae of *Anopheles hermsi* Barr. In choice experiments, larvae also selected high density patches of *M. aquaticum* stems as microhabitat. Similarly, emergence of *Culex erythrothorax* Dyar was positively correlated with the density of emergent bulrush stems and larvae were virtually absent from open areas of water in an experimental wetland (Workman & Walton, 2000). Survival of *Anopheles* larvae in the presence of mosquito fish, *Gambusia affinis* (Baird and Girard), was dependent on the density of predators and the density of macrophyte stems available as refuge (Orr & Resh, 1989). In contrast, Blaustein and Byard (1993) observed no evidence that a submergent plant (*Ceratophyllum demersum* L.) provided a refuge from a fish predator for *Culex pipiens* L. larvae.

Aquatic arthropods can respond to the presence of predators by reducing time spent in foraging and feeding behaviours (Macchiusi & Baker, 1992; Peckarsky *et al.*, 1993), increasing their propensity to use water currents to drift away (horizontally or vertically) from potential predators (Peckarsky & Penton, 1989; Brancelj & Blejec, 1994; McIntosh *et al.*, 2002), and by modifying life-history traits to reduce their period of occupancy of a dangerous habitat (Stibor, 1992). When such experiments involve exposure only to water previously inhabited by predators, the causes of changes in aquatic arthropod behaviour and life-history traits can be assigned to specific anti-predation responses (Sih, 1986; Burks *et al.*, 2000; Peckarsky *et al.*, 2002), whereas evaluation of direct exposure to predation involves a combination of antipredator responses and the result of selection of survivors with possibly specific phenotypic characteristics; survivors represent those less vulnerable prey that managed to avoid predators (Hechtel & Juliano, 1997).

The presence of fish resulted in emergence of adult females with consistently shorter wings compared with those that developed in the no-fish treatments. There was an interaction with algal cover in that, in the 33% algal cover treatment, wing lengths at the density of four fish were slightly greater than those observed in the one and two fish densities. A similar pattern was observed between treatments in the wing lengths of adult males, indicating that these effects are not sex-specific, in contrast to the sexual differences in responses to predators reported by Ball and Baker (1996) in a chironomid.

The identification of food sources and their relative importance in the larval diet are particularly relevant to certain novel approaches to vector control. The use of

biologically derived insecticides (Federici *et al.*, 2003; Bond *et al.*, 2004a), encapsulation of bacterial toxins in mosquito food items (Manasherob *et al.*, 1997), genetic modification of potential food plants (Wu *et al.*, 1997), and habitat manipulation (Knight *et al.*, 2003; Bond *et al.*, 2004b; Jiannino & Walton, 2004), all depend on knowledge of mosquito oviposition habitat selection and the ensuing consequences for larval diet selection. In conclusion, *An. pseudopunctipennis* selects oviposition habitats to ensure a suitable larval food source, filamentous algae. The intricate architecture of these algae represent a physical barrier to predation and thereby act as an effective refuge from key predators, such as *P. sphenops*.

### Acknowledgements

We thank Javier Valle, Rene Monzon, Eufonio Diaz, Pedro García, José L. Espinosa, and Oscar Reyes for technical help and Cuauhtémoc Villarreal for supplying mosquito larvae. This study received financial support from CONACYT-SIBEJ CHIS-S008. T.W. received financial assistance from the Plan Tecnológico de Navarra (UPNA) travel fund.

### References

- Angelon, K.A. & Petranka, J.W. (2002) Chemicals of predatory mosquitofish (*Gambusia affinis*) influence selection of oviposition site by *Culex* mosquitoes. *Journal of Chemical Ecology*, **28**, 797–806.
- Armbruster, P. & Hutchinson, R. (2002) Pupal mass and wing length as indicators of fecundity in *Aedes albopictus* and *Aedes geniculatus* (Diptera: Culicidae). *Journal of Medical Entomology*, **39**, 699–704.
- Ault, S.K. (1994) Environmental management: a re-emerging vector control strategy. *American Journal of Tropical Medicine and Hygiene*, **50**, 35–49.
- Ball, S.L. & Baker, R.L. (1996) Predator-induced life history changes: antipredator behavior costs or facultative life history shifts? *Ecology*, **77**, 1116–1124.
- Bentley, M.D. & Day, J.F. (1989) Chemical ecology and behavioral aspects of mosquito oviposition. *Annual Review of Entomology*, **34**, 401–421.
- Bentley, M.D., McDaniel, I.N., Lee, H.P., Stiehl, B. & Yatagai, M. (1976) Studies of *Aedes triseriatus* oviposition attractants produced by larvae of *Aedes triseriatus* and *Aedes atropalpus*. *Journal of Medical Entomology*, **13**, 112–115.
- Blaustein, L. (1999) Oviposition site selection in response to risk of predation: evidence from aquatic habitats and consequences for population dynamics and community structure. *Evolutionary Theory and Processes: Modern Perspectives* (ed. by S. P. Wasser), pp. 441–456. Kluwer, Dordrecht, Netherlands.
- Blaustein, L. & Byard, R. (1993) Predation by a cyprinodontid fish, *Aphanius mento*, on *Culex pipiens*: effects of alternative prey and vegetation. *Journal of the American Mosquito Control Association*, **9**, 356–358.
- Blaustein, L., Kiflawi, M., Eitam, A., Mangel, M. & Cohen, J.E. (2004) Oviposition habitat selection in response to risk of

- predation in temporary pools: mode of detection and consistency across experimental venue. *Oecologia*, **138**, 300–305.
- Blaustein, L. & Kotler, B.P. (1993) Oviposition habitat selection by the mosquito *Culiseta longiareolata*: effects of conspecifics, food, and green toad tadpoles. *Ecological Entomology*, **18**, 104–108.
- Bond, J.G., Marina, C.F. & Williams, T. (2004a) The naturally-derived insecticide spinosad is highly toxic to *Aedes* and *Anopheles* mosquito larvae. *Medical and Veterinary Entomology*, **18**, 50–56.
- Bond, J.G., Rojas, J.C., Arredondo-Jiménez, J.I., Quiroz-Martínez, H., Valle, J. & Williams, T. (2004b) Population control of the malaria vector *Anopheles pseudopunctipennis* by habitat manipulation. *Proceedings of the Royal Society London Series B, Biological Sciences*, **271**, 2161–2169.
- Brancelj, A. & Blejec, A. (1994) Diurnal vertical migration of *Daphnia hyaline* Leydig, 1860 (Crustacea: Cladocera) in Lake Bled (Slovenia) in relation to temperature and predation. *Hydrobiologia*, **284**, 125–136.
- Briegel, H. (1990a) Fecundity, metabolism, and body size in *Anopheles* (Diptera: Culicidae), vectors of malaria. *Journal of Medical Entomology*, **27**, 839–850.
- Briegel, H. (1990b) Metabolic relationship between female body size, reserves, and fecundity of *Aedes aegypti*. *Journal of Insect Physiology*, **36**, 165–172.
- Buma, A.G.J., Boelen, P. & Jeffrey, W.A. (2003) UVR-induced DNA damage in aquatic organisms. *UV Effects in Aquatic Organisms and Ecosystems* (ed. by W. Helbling and H. Zagarese), pp. 291–328. Royal Society of Chemistry, London.
- Burks, R.L., Jeppesen, E. & Lodge, D.M. (2000) Macrophyte and fish chemicals suppress *Daphnia* growth and alter life-history traits. *Oikos*, **88**, 139–147.
- Clements, A.N. (1992) *The Biology of Mosquitoes*. Vol. 1. *Development, Nutrition and Reproduction*. Chapman & Hall, London.
- Donahue, W.F. & Schindler, D.W. (1998) Diel emigration and colonization responses of blackfly larvae (Diptera: Simuliidae) to ultraviolet radiation. *Freshwater Biology*, **40**, 357–365.
- Eitam, A. & Blaustein, L. (2004) Oviposition habitat selection by mosquitoes in response to predator (*Notonecta maculata*) density. *Physiological Entomology*, **29**, 188–191.
- Federici, B.A., Park, H.Y., Bideshi, D.K., Wirth, M.C. & Johnson, J.J. (2003) Recombinant bacteria for mosquito control. *Journal of Experimental Biology*, **206**, 3877–3885.
- Fernández-Salas, I., Roberts, D.R., Rodríguez, M.H. & Marina-Fernández, C.F. (1994a) Bionomics of larval populations of *Anopheles pseudopunctipennis* in the Tapachula foothills area, southern Mexico. *Journal of the American Mosquito Control Association*, **10**, 477–486.
- Fernández-Salas, I., Rodríguez, M.H., Roberts, D.R., Rodríguez, M.C. & Wirtz, R.A. (1994b) Bionomics of adult *Anopheles pseudopunctipennis* (Diptera: Culicidae) in the Tapachula foothills area of southern Mexico. *Journal of Medical Entomology*, **31**, 663–670.
- Hall, T.F. (1972) The influence of plants on anopheline breeding. *American Journal of Tropical Medicine and Hygiene*, **21**, 787–794.
- Hechtel, L.J. & Juliano, S.A. (1997) Effects of a predator on prey metamorphosis: plastic responses by prey or selective mortality? *Ecology*, **78**, 838–851.
- Howland, L.J. (1930) Bionomical investigation of English mosquito larvae with special reference to their algal food. *Journal of Ecology*, **18**, 81–125.
- Jiannino, J.A. & Walton, W.E. (2004) Evaluation of vegetation management strategies for controlling mosquitoes in a southern California constructed wetland. *Journal of the American Mosquito Control Association*, **20**, 18–26.
- Kant, R., Pandey, S.D. & Sharma, S.K. (1996) Mosquito breeding in relation to aquatic vegetation and some physico-chemical parameters in rice fields of central Gujarat. *Indian Journal of Malariology*, **33**, 30–40.
- Keiper, J.B. (2002) Biology and immature stages of coexisting Hydroptilidae (Trichoptera) from Northeastern Ohio lakes. *Annals of the Entomological Society of America*, **95**, 608–616.
- Kiflawi, M., Blaustein, L. & Mangel, M. (2003) Oviposition habitat selection by the mosquito *Culiseta longiareolata* in response to risk of predation and conspecific larval density. *Ecological Entomology*, **28**, 168–173.
- Knight, R.L., Walton, W.E., O'Meara, G.F., Reisen, W.K. & Wass, R. (2003) Strategies for effective mosquito control in constructed treatment wetlands. *Ecological Engineering*, **21**, 211–232.
- Laird, M. (1988) *The Natural History of Larval Mosquito Habitats*. Academic Press, London.
- Laurence, B.R. & Pickett, J.A. (1985) An oviposition attractant pheromone in *Culex quinquefasciatus* Say (Diptera: Culicidae). *Bulletin of Entomological Research*, **75**, 283–290.
- Macchiusi, F. & Baker, R.L. (1992) Effects of predators and food availability on activity and growth of *Chironomus tentans* (Chironomidae: Diptera). *Freshwater Biology*, **28**, 207–216.
- Manasherob, R., Bendov, E., Margalit, J., Zaritsky, A. & Barak, Z. (1997) Raising activity of *Bacillus thuringiensis* var. *israelensis* against *Anopheles stephensi* larvae by encapsulation in *Tetrahymena pyriformis* (Hymenostomatida, Tetrahymenidae). *Journal of the American Mosquito Control Association*, **12**, 627–631.
- Manatunge, J., Asaeda, T. & Priyadarshana, T. (2000) The influence of structural complexity on fish-zooplankton interactions: a study using artificial submerged macrophytes. *Environmental Biology of Fishes*, **58**, 425–438.
- Manguin, S., Roberts, D.R., Peyton, E.L., Rejmankova, E. & Pecor, J. (1996) Characterization of *Anopheles pseudopunctipennis* larval habitats. *Journal of the American Mosquito Control Association*, **12**, 619–626.
- Marten, G.G., Suarez, M.F. & Astaeza, R. (1996) An ecological survey of *Anopheles albimanus* larval habitats in Colombia. *Journal of Vector Ecology*, **21**, 122–131.
- Mboera, L.E.G., Mdira, K.Y., Salum, F.M., Takken, W. & Pickett, J.A. (1999) Influence of synthetic oviposition pheromone and volatiles from soakage pits and grass infusions upon oviposition site-selection of *Culex* mosquitoes in Tanzania. *Journal of Chemical Ecology*, **25**, 1855–1865.
- McIntosh, A.R., Peckarsky, B.L. & Taylor, B.W. (2002) The influence of predatory fish on mayfly drift: exploiting from experiments to nature. *Freshwater Biology*, **47**, 1497–1513.
- Merritt, R.W., Dadd, R.H. & Walker, E.D. (1992) Feeding behavior, natural food, and nutritional relationships of larval mosquitoes. *Annual Review of Entomology*, **37**, 349–376.
- Merritt, R.W., Olds, E.J. & Walker, E.D. (1990) Natural food and feeding behavior of *Coquillettidia perturbans* larvae. *Journal of the American Mosquito Control Association*, **6**, 35–41.
- Orr, B.K. & Resh, V.H. (1989) Experimental test of the influence of aquatic macrophyte cover on the survival of *Anopheles* larvae. *Journal of the American Mosquito Control Association*, **5**, 579–585.
- Orr, B.K. & Resh, V.H. (1992) Influence of *Myriophyllum aquaticum* cover on *Anopheles* mosquito abundance, oviposition, and larval microhabitat. *Oecologia*, **90**, 474–482.
- Peckarsky, B.L., Cowan, C.A., Penton, M.A. & Anderson, C. (1993) Sublethal consequences of stream-dwelling predatory



- stoneflies on mayfly growth and fecundity. *Ecology*, **74**, 1836–1846.
- Peckarsky, B.L., McIntosh, A.R., Taylor, B.W. & Dahl, J. (2002) Predatory chemicals induce changes in mayfly life history traits: a whole stream manipulation. *Ecology*, **83**, 612–618.
- Peckarsky, B.L. & Penton, M.A. (1989) Mechanisms of prey selection by stream-dwelling stoneflies. *Ecology*, **70**, 1203–1218.
- Peckarsky, B.L., Taylor, B.W. & Caudill, C.C. (2000) Hydrologic and behavioral constraints on oviposition of stream insects: implications for adult dispersal. *Oecologia*, **125**, 186–200.
- Persson, L. & Eklov, P. (1995) Prey refuges affecting interactions between piscivorous perch and juvenile perch and roach. *Ecology*, **76**, 70–81.
- Reiskind, M.H. & Wilson, M.L. (2004) *Culex restuans* (Diptera: Culicidae) oviposition behavior determined by larval habitat quality and quantity in southeastern Michigan. *Journal of Medical Entomology*, **41**, 179–186.
- Rejmankova, E., Harbin-Ireland, A. & Lege, M. (2000) Bacterial abundance in larval habitats of four species of *Anopheles* (Diptera: Culicidae) in Belize, Central America. *Journal of Vector Ecology*, **25**, 229–238.
- Rejmankova, E., Roberts, D.R., Harbach, R.E., Pecor, J., Peyton, E.L., Manguin, S. *et al.* (1993) Environmental and regional determinants of *Anopheles* (Dipt: Culicidae) larval distribution in Belize, Central America. *Environmental Entomology*, **22**, 978–992.
- Rejmankova, E.H., Savage, M., Rejmanek, M., Arredondo-Jiménez, J.I. & Roberts, D.R. (1991) Multivariate analysis of relationships between habitats, environmental factors and occurrence of anopheline mosquito larvae *Anopheles albimanus* and *An. pseudopunctipennis* in southern Chiapas, Mexico. *Journal of Applied Ecology*, **28**, 827–841.
- Rejmankova, E.H., Savage, M., Roberts, D.R., Manguin, S., Pope, K.O., Komárek, J. *et al.* (1996) *Anopheles albimanus* (Diptera: Culicidae) and cyanobacteria: an example of larval habitat selection. *Environmental Entomology*, **25**, 1058–1087.
- Rejmankova, E., Savage, H.M., Rodríguez, M.H., Roberts, D.R. & Rejmanek, M. (1992) Aquatic vegetation as a basis or classification of *Anopheles albimanus* Weidemann (Diptera: Culicidae) larval habitats. *Environmental Entomology*, **21**, 598–603.
- Savino, J.F. & Stein, R.A. (1989) Behavioural interactions between fish predators and their prey: effects of plant density. *Animal Behaviour*, **37**, 311–321.
- Service, M.W. (1989) The importance of ecological studies on malaria vectors. *Bulletin of the Society for Vector Ecology*, **14**, 26–38.
- Service, M.W. (1993) *Mosquito Ecology. Field Sampling Methods*, 2nd edn. Elsevier, London.
- Service, M.W. (1995) Can we control mosquitoes without pesticides? A summary. *Journal of the American Mosquito Control Association*, **11**, 290–293.
- Sih, A. (1986) Antipredatory responses and the perception of danger by mosquito larvae. *Ecology*, **67**, 434–441.
- Sokal, R.R. & Rohlf, F.J. (1981) *Biometry*, 2nd edn. Freeman, New York.
- Spencer, M., Blaustein, L. & Cohen, J.E. (2002) Oviposition habitat selection by mosquitoes (*Culiseta longiareolata*) and consequences for population size. *Ecology*, **83**, 669–679.
- SPSS (1995) *SPSS for Windows, Version 10.0*. SPSS Science Inc., Chicago, Illinois.
- Stav, G., Blaustein, L. & Margalith, J. (1999) Experimental evidence for predation risk sensitive oviposition by a mosquito, *Culiseta longiareolata*. *Ecological Entomology*, **24**, 202–207.
- Stibor, H. (1992) Predator induced life-history shifts in a freshwater cladoceran. *Oecologia*, **92**, 162–165.
- Villarreal, C., Arredondo-Jiménez, J.I., Rodríguez, M.H. & Ulloa, A. (1998) Colonization of *Anopheles pseudopunctipennis* from Mexico. *Journal of Medical Entomology*, **14**, 369–372.
- Walker, E.D., Olds, E.J. & Merritt, R.W. (1988) Gut content analysis of mosquito larvae (Diptera: Culicidae) using DAPI stain and epifluorescence microscopy. *Journal of Medical Entomology*, **25**, 351–354.
- Wetzel, R.G. & Likens, G.E. (1991) *Limnological Analyses*, 2nd edn. Springer-Verlag, New York.
- Workman, P.D. & Walton, W.E. (2000) Emergence patterns of *Culex* mosquitoes (Diptera: Culicidae) at an experimental constructed treatment wetland in southern California. *Journal of the American Mosquito Control Association*, **16**, 124–130.
- Wotton, R.S., ed. (1990) *The Biology of Particles in Aquatic Systems*. CRC Press, Boca Raton, Florida.
- Wu, X., Vennison, S.J., Huirong, L., Ben-Dov, E., Zaritsky, A. & Boussiba, S. (1997) Mosquito larvicidal activity of transgenic *Anabaena* strain PCC 7120 expressing combinations of genes from *Bacillus thuringiensis* subsp. *israelensis*. *Applied and Environmental Microbiology*, **63**, 4971–4974.
- Zack, R.S. & Foote, B.A. (1978) Utilization of algal monocultures by larvae of *Scatella stagnalis*. *Environmental Entomology*, **7**, 509–511.
- Zahiri, N. & Rau, M.E. (1998) Oviposition attraction and repellency of *Aedes aegypti* (Diptera: Culicidae) to waters from conspecific larvae subjected to crowding, confinement, starvation or infection. *Journal of Medical Entomology*, **35**, 782–787.

Accepted 7 December 2004